

WHAT IS CLAIMED IS:

Sub
Q1
5 1. A pharmacologically active peptide conjugate having a reduced tendency towards enzymatic cleavage comprising X and Z,

wherein X is a pharmacologically active peptide sequence, and

wherein Z is a stabilising peptide sequence, of 4-20 amino acid units covalently bound to X, wherein each amino acid unit in said stabilising peptide sequence, Z is selected from the
10 group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I



15 wherein R¹ and R² are selected from the group consisting of hydrogen, C₁₋₆-alkyl, phenyl, and phenyl-methyl, wherein C₁₋₆-alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and
20 carboxy, or R¹ and R² together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring, e.g. 2,4-diaminobutanoic acid and 2,3-diaminopropanoic acid; and

wherein the ratio between the half-life of said peptide conjugate and the half-life of the
25 corresponding pharmacologically active peptide sequence, X, when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least about 2, preferably at least about 3, such as at least about 5, more preferably at least about 7, such as at least about 9, e.g. at least about 10 or when said pharmacologically active peptide X is not orally
30 absorbed, said conjugate is absorbed,

05341590-021399

or a salt thereof, with the proviso that said pharmacologically active peptide conjugate is not selected from

H-Trp-Ala-Gly-Gly-Asp-Ala-Ser-Gly-Glu-(Lys-Glu)₃-OH,

H-Trp-Ala-Gly-Gly-Asp-Ala-Ser-Gly-Glu-(Glu)₆-OH,

5 H-Tyr-Gly-Gly-Phe-Leu-(Glu)₆-OH and

H-Tyr-Gly-Gly-Phe-Leu-(Lys)₆-OH.

2. A peptide conjugate according to claim 1, wherein Z is covalently bound to X at the C-terminal carbonyl function of X.

10

3. A peptide conjugate according to claim 1, wherein Z is covalently bound to the N-terminal nitrogen atom of X.

4. A peptide conjugate according to claim 1, wherein the first sequence (Z) is covalently
15 bound to X at the C-terminal carbonyl function of X and the second sequence (Z) is covalently bound to the N-terminal nitrogen atom of X.

5. A peptide conjugate according to claim 1, wherein Z is covalently bound to a nitrogen atom on the side chain of a lysine, arginine or histidine residue or a carbonyl function on the
20 side chain of glutamic acid or aspartic acid of X.

6. A peptide conjugate according to ^{Claim 1} ~~any of the preceding claims~~, wherein Z consists of 4-15, preferably 4-10, more preferably 4-7, such as 6 amino acid units.

7. A peptide conjugate according to claim ~~5 or 6~~ ¹, wherein each amino acid unit in Z is independently selected from the group consisting of Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Orn, 2,4-diaminobutanoic acid, 2,3-diaminopropanoic acid and Met.

8. A peptide conjugate according to claim 7, wherein each amino acid unit in Z is
30 selected from the group consisting of Glu, Lys and Met.

0944590.01389

sub
C2

a

a sub C3

87
clean 1
any of clo

9. A peptide conjugate according to ~~any of claims 5 to 8~~, wherein Z comprises at least one Lys amino acid unit, preferably at least two Lys amino acid units, such as at least three Lys amino acid units, e.g. at least four Lys amino acid units, more preferably at least five Lys amino acid units, such as six Lys amino acid units.

5

10. A peptide conjugate according to claim 9, wherein Z is (Lys)_n, wherein n is an integer in the range from 4 to 15, preferably in the range from 4 to 10, such as in the range from 4 to 8, e.g. in the range from 4 to 6.

Sub
D1

10 11.

12.

- 10 11.) A peptide conjugate according to claim 9, wherein Z is Lys₄, Lys₅ or Lys₆.

12. A peptide conjugate according to claim 11, wherein Z is Lys₆.

claim 1
any of cl

13. A peptide conjugate according to any of claims 5 to 9, wherein Z is (Lys-Xaa)_m or
15 (Xaa-Lys)_m, wherein m is an integer in the range from 2 to 7, and each Xaa is independently
selected from the group consisting of Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Arg, His, Orn, 2,4-
diaminobutanoic acid, 2,3-diaminopropanoic acid and Met.

14. A peptide conjugate according to claim 13, wherein Z is (Lys-Xaa)₃ or (Xaa-Lys)₃,
20 wherein each Xaa is independently selected from the group consisting of Ser, Thr, Tyr, Asn,
Gln, Asp, Glu, Arg, His, Orn, 2,4-diaminobutanoic acid, 2,3-diaminopropanoic acid and
Met.

Sub
25

15. A peptide conjugate according to claim 15, wherein Z is (Lys-Glu)₃ or (Glu-Lys)₃.

claim 1
any of

16. A peptide conjugate according to ~~any of claims 5 to 9~~, wherein Z is Lys_p-Xaa_q or Xaa_p-Lys_q, wherein p and q are integers in the range from 1 to 14, with the proviso that p+q is in the range of 3-15, and each Xaa is independently selected from the group consisting of Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Arg, His, Orn, 2,4-diaminobutanoic acid, 2,3-diaminopropanoic acid and Met.

17. A peptide conjugate according to claim 16, wherein Z is Lys₃-Xaa₃ or Xaa₃-Lys₃, wherein each Xaa is independently selected from the group consisting of Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Arg, His, Orn, 2,4-diaminobutanoic acid, 2,3-diaminopropanoic acid and Met.

18. A peptide conjugate according to claim 17, wherein Z is Lys₃-Glu₃ or Glu₃-Lys₃,

19. A peptide conjugate according to claims 5-18, wherein Z is Lys-Lys-Lys-Lys, Xaa-Lys-Lys-Lys, Lys-Xaa-Lys-Lys, Lys-Lys-Xaa-Lys, Lys-Lys-Lys-Xaa, Xaa-Xaa-Lys-Lys, Xaa-

10 Lys-Xaa-Lys, Xaa-Lys-Lys-Xaa, Lys-Xaa-Xaa-Lys, Lys-Xaa-Lys-Xaa, Lys-Lys-Xaa-Xaa,
Xaa-Xaa-Xaa-Lys, Xaa-Xaa-Lys-Xaa, Xaa-Lys-Xaa-Xaa, Lys-Xaa-Xaa-Xaa, Xaa-Xaa-
Xaa-Xaa, Lys-Lys-Lys-Lys-Lys, Xaa-Lys-Lys-Lys-Lys, Lys-Xaa-Lys-Lys-Lys, Lys-Lys-
Xaa-Lys-Lys, Lys-Lys-Lys-Xaa-Lys, Lys-Lys-Lys-Lys-Xaa, Xaa-Xaa-Lys-Lys-Lys, Xaa-
Lys-Xaa-Lys-Lys, Xaa-Lys-Lys-Xaa-Lys, Xaa-Lys-Lys-Lys-Xaa, Lys-Xaa-Xaa-Lys-Lys,
15 Lys-Xaa-Lys-Xaa-Lys, Lys-Xaa-Lys-Lys-Xaa, Lys-Lys-Xaa-Xaa-Lys, Lys-Lys-Xaa-Lys-
Xaa, Lys-Lys-Lys-Xaa-Xaa, Lys-Lys-Xaa-Xaa-Xaa, Lys-Xaa-Lys-Xaa-Xaa, Lys-Xaa-Xaa-
Lys-Xaa, Lys-Xaa-Xaa-Xaa-Lys, Xaa-Lys-Lys-Xaa-Xaa, Xaa-Lys-Xaa-Xaa-Lys, Xaa-Xaa-
Lys-Lys-Xaa, Xaa-Xaa-Lys-Xaa-Lys, Xaa-Xaa-Xaa-Lys-Lys, Lys-Xaa-Xaa-Xaa-Xaa, Xaa-
Lys-Xaa-Xaa-Xaa, Xaa-Xaa-Lys-Xaa-Xaa, Xaa-Xaa-Xaa-Lys-Xaa, Xaa-Xaa-Xaa-Xaa-Lys,
20 Xaa-Xaa-Xaa-Xaa-Xaa, Lys-Lys-Lys-Lys-Lys-Lys, Xaa-Lys-Lys-Lys-Lys-Lys, Lys-Xaa-
Lys-Lys-Lys-Lys, Lys-Lys-Xaa-Lys-Lys-Lys, Lys-Lys-Lys-Xaa-Lys-Lys, Lys-Lys-Lys-
Lys-Xaa-Lys, Lys-Lys-Lys-Lys-Lys-Xaa, Xaa-Xaa-Lys-Lys-Lys-Lys, Xaa-Lys-Xaa-Lys-
Lys-Lys, Xaa-Lys-Lys-Xaa-Lys-Lys, Xaa-Lys-Lys-Lys-Xaa-Lys, Xaa-Lys-Lys-Lys-Lys-
Xaa, Lys-Xaa-Xaa-Lys-Lys-Lys, Lys-Xaa-Lys-Xaa-Lys-Lys, Lys-Xaa-Lys-Lys-Xaa-Lys,
25 Lys-Xaa-Lys-Lys-Lys-Xaa, Lys-Lys-Xaa-Xaa-Lys-Lys, Lys-Lys-Xaa-Lys-Xaa-Lys, Lys-
Lys-Xaa-Lys-Lys-Xaa, Lys-Lys-Lys-Xaa-Xaa-Lys, Lys-Lys-Lys-Xaa-Lys-Xaa, Lys-Lys-
Lys-Lys-Xaa-Xaa, Xaa-Xaa-Xaa-Lys-Lys-Lys, Xaa-Xaa-Lys-Xaa-Lys-Lys, Xaa-Xaa-Lys-
Lys-Xaa-Lys, Xaa-Xaa-Lys-Lys-Lys-Xaa, Xaa-Lys-Xaa-Xaa-Lys-Lys, Xaa-Lys-Xaa-Lys-
Xaa-Lys, Xaa-Lys-Xaa-Lys-Lys-Xaa, Xaa-Lys-Lys-Xaa-Xaa-Lys, Xaa-Lys-Lys-Xaa-Lys-
30 Xaa, Xaa-Lys-Lys-Lys-Xaa-Xaa, Lys-Lys-Lys-Xaa-Xaa-Xaa, Lys-Lys-Xaa-Lys-Xaa-Xaa,
Lys-Lys-Xaa-Xaa-Lys-Xaa, Lys-Lys-Xaa-Xaa-Xaa-Lys, Lys-Xaa-Lys-Lys-Xaa-Xaa, Lys-
Xaa-Lys-Xaa-Lys-Xaa, Lys-Xaa-Lys-Xaa-Xaa-Lys, Lys-Xaa-Xaa-Lys-Lys-Xaa, Lys-Xaa-

090705Z JUL 68

- Xaa-Lys-Xaa-Lys, Lys-Xaa-Xaa-Xaa-Lys-Lys, Lys-Lys-Xaa-Xaa-Xaa-Xaa, Lys-Xaa-Lys-Xaa-Xaa-Xaa, Lys-Xaa-Xaa-Lys-Xaa-Xaa, Lys-Xaa-Xaa-Xaa-Lys-Xaa, Lys-Xaa-Xaa-Xaa-Xaa-Lys, Xaa-Lys-Lys-Xaa-Xaa-Xaa, Xaa-Lys-Xaa-Lys-Xaa-Xaa, Xaa-Lys-Xaa-Xaa-Lys-Xaa, Xaa-Lys-Xaa-Xaa-Xaa-Lys, Xaa-Xaa-Lys-Lys-Xaa-Xaa, Xaa-Xaa-Lys-Xaa-Lys-Xaa, Xaa-Xaa-Lys-Xaa-Xaa-Lys, Xaa-Xaa-Xaa-Lys-Lys-Xaa, Xaa-Xaa-Xaa-Lys-Lys-Lys, Xaa-Xaa-Xaa-Lys-Xaa-Lys, Xaa-Xaa-Xaa-Xaa-Lys-Lys, Lys-Xaa-Xaa-Xaa-Xaa-Xaa, Xaa-Lys-Xaa-Xaa-Xaa-Xaa, Xaa-Xaa-Lys-Xaa-Xaa-Xaa, Xaa-Xaa-Xaa-Lys-Xaa-Xaa, Xaa-Xaa-Xaa-Lys-Xaa, Xaa-Xaa-Xaa-Xaa-Xaa-Lys, or Xaa-Xaa-Xaa-Xaa-Xaa-Xaa, wherein each Xaa is independently selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Arg, His, Met, Orn, and amino acids of the formula I



- wherein R¹ and R² are selected from the group consisting of hydrogen, C₁₋₆-alkyl, phenyl, and phenyl-methyl, wherein C₁₋₆-alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R¹ and R² together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring, e.g. 2,4-diaminobutanoic acid (Dbu) and 2,3-diaminopropanoic acid (Dpr).

20. A peptide conjugate according to ~~any of claims 5 to 18~~, wherein Z consists of L-amino acids only.

21. A peptide conjugate according to ~~any of claims 5-7~~, wherein Z is (Dbu)_n or (Dpr)_n, wherein n is an integer in the range from 4 to 15, preferably in the range from 4 to 10, such as in the range from 4 to 8, e.g. in the range from 4 to 6.

22. A peptide conjugate according to claim 1, wherein Z is Dpr₆.

034150-01399

a

svb
c 4

25

claim 1

claim 1

Sub C5
23. A peptide conjugate according to any of the preceding claims, wherein said pharmacologically active peptide sequence (X) consists of at the most 75 amino acid units, such as at the most 65, e.g. at the most 60, preferably at the most 55, such as at the most 53, e.g. at the most 50.

Sub B4
5- 24. The peptide conjugate according to claim 16, wherein X is selected from the group consisting of enkephalin, Leu-enkephalin, Met-enkephalin, angiotensin I, angiotensin II, vasopressin, endothelin, vasoactive intestinal peptide, neurotensin, endorphins, insulin, gramicidin, paracelsin, delta-sleep inducing peptide, gonadotropin-Releasing hormone,
10 human parathyroid hormone (1-34), truncated erythropoietin analogues, specifically EMP-1, Atrial natriuretic peptide (ANP, ANF), human brain natriuretic peptide (hBNP), cecropin, kinetensin, neurophysins, elafin, guamerin, atriopeptin I, atriopeptin II, atriopeptin III, deltorphin I, deltorphin II, vasotocin, bradykinin, dynorphin, dynorphin A, dynorphin B, growth hormone release factor, growth hormone, growth hormone releasing peptide,
15 oxytocin, calcitonin, calcitonin gene-related peptide, calcitonin gene-related peptide II, growth hormone releasing peptide, tachykinin, adrenocorticotrophic hormone (ACTH), brain natriuretic polypeptide, cholecystokinin, corticotropin releasing factor, diazepam binding inhibitor fragment, FMRF-amide, galanin, gastric releasing polypeptide, gastric inhibitory polypeptide, gastrin, gastrin releasing peptide, glucagon, glucagon-like peptide-1, glucagon-
20 like peptide-2, LHRH, melanin concentrating hormone, melanocyte stimulating hormone (MSH), alpha-MSH, morphine modulating peptides, motilin, neurokinin A, neurokinin B, neuromedin B, neuromedin C, neuromedin K, neuromedin N, neuromedin U, neuropeptide K, neuropeptide Y, pituitary adenylate cyclase activating polypeptide (PACAP), pancreatic polypeptide, peptide YY, peptide histidine-methionine amide (PHM), secretin,
25 somatostatin, substance K, thyrotropin-releasing hormone (TRH), kyotorphin, melanostatin (MIF-1), thrombopoietin analogs, in particular AF 12505 (Ile-Glu-Gly-Pro-Thr-Leu-Arg-Gln-Trp-Leu-Ala-Ala-Arg-Ala), insulin-like growth factor I (57-70) (Ala-Leu-Leu-Glu-Thr-Tyr-Cys-Ala-Thr-Pro-Ala-Lys-Ser-Glu), insulin-like growth factor I (30-41) (Gly-Tyr-Gly-Ser-Ser-Ser-Arg-Arg-Ala-Pro-Gln-Thr), insulin-like growth factor I (24-41) (Tyr-Phe-
30 Asn-Lys-Pro-Thr-Gly-Tyr-Gly-Ser-Ser-Ser-Arg-Arg-Ala-Pro-Gln-Thr), insulin-like growth factor II (33-40) (Ser-Arg-Val-Ser-Arg-Arg-Ser-Arg), insulin-like growth [tyro] factor II (33-40) (Tyr-Ser-Arg-Val-Ser-Arg-Arg-Ser-Arg), insulin-like growth factor II (69-84)

05341590-071399

(Asp-Val-Ser-Thr-Pro-Pro-Thr-Val-Leu-Pro-Asp-Asn-Phe-Pro-Arg-Tyr), growth hormone (GH)-releasing peptide-6 (GHRP-6) (His-DTrp-Ala-Trp-DPhe-Lys-NH₂), beta-Interleukin I (163-171) (Val-Gln-Gly-Glu-Glu-Ser-Asn-Asp-Lys), beta-Interleukin II (44-56) (Ile-Leu-Asn-Gly-Ile-Asn-Asn-Tyr-Lys-Asn-Pro-Lys-Leu), Interleukin II (60-70) (Leu-Thr-Phe-Lys-Phe-Tyr-Met-Pro-Lys-Lys-Ala), exendin-4 (GLP-1 analog) (His-Gly-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Leu-Ser-Lys-Gln-Met-Glu-Glu-Glu-Ala-Val-Arg-Leu-Phe-Ile-Glu-Trp-Leu-Lys-Asn-Gly-Gly-Pro-Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser-NH₂), exendin-3 (GLP-1 analog) (His-Ser-Asp-Gly-Thr-Phe-Thr-Ser-Asp-Leu-Ser-Lys-Gln-Met-Glu-Glu-Glu-Ala-Val-Arg-Leu-Phe-Ile-Glu-Trp-Leu-Lys-Asn-Gly-Gly-Pro-Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser), [Cys(Acm)_{20,31}] epidermal growth factor (20-31) Cys(Acm)-Met-His-Ile-Glu-Ser-Leu-Asp-Ser-Tyr-Thr-Cys(Acm), bivalirudin (Hirulog) (D-Phe-Pro-Arg-Pro-(Gly)₄-Asn-Gly-Asp-Phe-Glu-Glu-Ile-Pro-Glu-Glu-Tyr-Leu), hirulog-1 D-Phe-Pro-Arg-Pro-(Gly)₄-Asn-Gly-Asp-Phe-Glu-Glu-Ile-Pro-Glu-Tyr-Leu, C-type natriuretic peptide (1-53) (CNP) (Asp-Leu-Arg-Val-Asp-Thr-Lys-Ser-Arg-Ala-Ala-Trp-Ala-Arg-Leu-Leu-Gln-Glu-His-Pro-Asn-Ala-Arg-Lys-Tyr-Lys-Gly-Ala-Asn-Lys-Lys-Gly-Leu-Ser-Lys-Gly-Cys-Phe-Gly-Leu-Lys-Leu-Asp-Arg-Ile-Gly-Ser-Met-Ser-Gly-Leu-Gly-Cys; Disulfide bridge: Cys₃₇-Cys₅₃), "Mini ANP" (Met-Cys-His-cyclohexylAla-Gly-Gly-Arg-Met-Asp-Arg-Ile-Ser-Cys-Tyr-Arg, disulfide bridge cys₂-cys₁₃), Melanotan-II (also known as MT-II, alpha-MSH₄₋₁₀-NH₂, or Ac-Nle₄-Asp₅-His₆-D-Phe₇-Arg₈-Trp₉-Lys₁₀), thymosin alpha₁ (TA₁) (Ac-Ser-Asp-Ala-Ala-Val-Asp-Thr-Ser-Ser-Glu-Ile-Thr-Thr-Lys-Asp-Leu-Lys-Glu-Lys-Lys-Glu-Val-Val-Glu-Glu-Ala-Glu-Asn), ornipressin (also known as 8-ornithine-vasopressin, (POR-8), [Phe₂,Ile₃,Orn₈]vasopressin), Cys-Phe-Ile-Gln-Asn-Cys-Pro-Orn-Gly-NH₂, Disulfide bridge: Cys₁-Cys₆), octreotide (201-995) (DPhe-Cys-Phe-DTrp-Lys-Thr-Cys-Thr-ol; disulfide bridge: Cys₂-Cys₇), eptifibatide (INTEGRILIN), calcitonin gene-related peptide (CGRP) (Ala-Cys-Asp-Thr-Ala-Thr-Cys-Val-Thr-His-Arg-Leu-Ala-Gly-Leu-Leu-Ser-Arg-Ser-Gly-Gly-Val-Val-Lys-Asn-Asn-Phe-Val-Pro-Thr-Asn-Val-Gly-Ser-Lys-Ala-Phe-NH₂; Disulfide bridge: Cys₂-Cys₇), endomorphin-1 Tyr-Pro-Trp-Phe-NH₂; endomorphin-2 Tyr-Pro-Phe-Phe-NH₂, nociceptin (also known as Orphanin FQ, Phe-Gly-Gly-Phe-Thr-Gly-Ala-Arg-Lys-Ser-Ala-Arg-Lys-Leu-Ala-Asn-Gln), angiotensinogen (1-13) (Asp-Arg-Val-Tyr-Ile-His-Pro-Phe-His-Leu-Val-Ile-His), adrenomodullin (1-12) (Tyr-Arg-Gln-Ser-Met-Asn-Asn-Phe-Gln-Gly-Leu-Arg), antiarrhythmic peptide (AAP) (Gly-Pro-Hyp-Gly-Ala-Gly), Antagonist G (Arg-DTrp-(nMe)Phe-DTrp-Leu-Met-NH₂), indolicidin

(Ile-Deu-Pro-Trp-Lys-Trp-Pro-Trp-Trp-Pro-Trp-Arg-Arg-NH₂), osteocalcin (37-49) (Gly-Phe-Gln-Glu-Ala-Tyr-Arg-Arg-Phe-Tyr-Gly-Pro-Val), cortistatin 29 (1-13) (Glp)-Glu-Arg-Pro-Pro-Leu-Gln-Gln-Pro-Pro-His-Arg-Asp), cortistatin 14 Pro-Cys-Lys-Asn-Phe-Phe-Trp-Lys-Thr-Phe-Ser-Ser-Cys-Lys; Disulfide bridge: Cys2-Cys13, PD-145065 (Ac-D-Bhg-Leu-Asp-Ile-Ile-Trp), PD-142893 (Ac-D-Dip-Leu-Asp-Ile-Ile-Trp), fibrinogen binding inhibitor peptide (His-His-Leu-Gly-Gly-Ala-Lys-Gln-Ala-Gly-Asp-Val), leptin (93-105) (Asn-Val-Ile-Gln-Ile-Ser-Asn-Asp-Leu-Glu-Asn-Leu-Arg), GR 83074 (Boc-Arg-Ala-DTrp-Phe-DPro-Pro-Nle-NH₂), Tyr-W-MIF-1 (Tyr-Pro-Trp-Gly-NH₂), parathyroid hormone related peptide (107-111) (Thr-Arg-Ser-Ala-Trp), angiotensinogen (1-14) Asp-Arg-Val-Tyr-Ile-His-Pro-Phe-His-Leu-Val-Ile-His-Asn, Leupeptin (Ac-Leu-Leu-Arg-CHO), and any modified or truncated analogue thereof.

25. A peptide conjugate according to ^{claim 1} any of the previous claims wherein the conjugate is

15 H-Tyr-Ala-Asp-Ala-Ile-Phe-Thr-Asn-Ser-Tyr-Arg-Lys-Val-Leu-Gly-Gln-Leu-Ser-Ala-Arg-Lys-Leu-Leu-Gln-Asp-Ile-Met-Ser Arg-Gln-Gln-Gly-Glu-Ser-Asn-Gln-Glu-Arg-Gly-Ala-Arg-Ala-Arg-Leu-Lys₆-NH₂ (GHRH(1-44)(Human)-Lys₆-NH₂);

H-Tyr-Ala-Asp-Ala-Ile-Phe-Thr-Asn-Ser-Tyr-Arg-Lys-Val-Leu-Gly-Gln-Leu-Ser-Ala-Arg-Lys-Leu-Leu-Gln-Asp-Ile-Met-Ser Arg-Gln-Gln-Gly-Glu-Ser-Asn-Gln-Glu-Arg-Gly-Ala-Arg-Ala-Arg-Leu-Glu₆-NH₂ (GHRH (1-44)(Human)-Glu₆-NH₂);

H- Lys₆-Ser-Val-Ser-Glu-Ile-Gln-Leu-Met-His-Asn-Leu-Gly-Lys-His-Leu-Asn-Ser-Met-Glu-Arg-Val-Glu-Trp-Leu-Arg-Lys-Lys-Leu-Gln-Asp-Val-His-Asn-Phe-OH (Lys₆-PTH(1-34)(Human)-OH);

H-Ser-Val-Ser-Glu-Ile-Gln-Leu-Met-His-Asn-Leu-Gly-Lys-His-Leu-Asn-Ser-Met-Glu-Arg-Val-Glu-Trp-Leu-Arg-Lys-Lys-Leu-Gln-Asp-Val-His-Asn-Phe-Lys₆-OH (PTH(1-34)(Human)-Lys₆-OH);

30 H-His-Ala-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-Ser-Tyr-Leu-Glu-Gly-Gln-Ala-Ala-Lys-Glu-Phe-Ile-Ala-Trp-Leu-Val-Lys-Gly-Arg-Lys₆-OH (GLP-1-(7-36)(Human)-Lys₆-OH);

66ET20"065T4E60

H-Gly-Gly-Thr-Tyr-Ser-Cys(Acm)-His-Phe-Gly-Pro-Leu-Thr-Trp-Val-Cys(Acm)-Lys-Pro-Gln-Gly-Gly-Lys₆-OH (EMP-1-Lys₆-OH);

5 H- Lys₆-Gly-Gly-Thr-Tyr-Ser-Cys(Acm)-His-Phe-Gly-Pro-Leu-Thr-Trp-Val-Cys(Acm)-Lys-Pro-Gln-Gly-Gly-OH (Lys₆-EMP-1-OH);

H- Lys₆-Gly-Gly-Thr-Tyr-Ser-Cys(Acm)-His-Phe-Gly-Pro-Leu-Thr-Trp-Val-Cys(Acm)-Lys-Pro-Gln-Gly-Gly-Lys₆-OH (Lys₆-EMP-1-Lys₆-OH);

10

H-Aib-His-2-D-Nal-D-Phe-Lys-(Lys)₆-NH₂ (GHRP-(Lys)₆-NH₂);

H-Tyr-Gly-Gly-Phe-Leu-Lys-Lys-Glu-Glu-Glu-Lys--OH (Leu-enkephalin-Lys-Lys-Glu-Glu-Glu-Lys-OH);

15

H-Tyr-Gly-Gly-Phe-Leu-Lys-Glu-Glu-Glu-Glu-Lys--OH (Leu-enkephalin-Lys-Glu-Glu-Glu-Glu-Lys-OH);

H-Tyr-Gly-Gly-Phe-Leu-Lys-Glu-Glu-Glu-Glu-Lys--OH (Leu-enkephalin-(Lys-Glu)₃);

20

H-Tyr-Gly-Gly-Phe-Leu-(Dpr)₆-OH (Leu-enkephalin-(Dpr)₆-OH);

H-Lys₆-Tyr-Gly-Gly-Phe-Leu-OH (H-Lys₆-Leu-enkephalin);

25 H-Tyr-Gly-Gly-Phe-Leu-Lys₆-OH (H-Leu-enkephalin-Lys₆);

H-Lys₆-Tyr-Gly-Gly-Phe-Leu-Lys₆-OH (H-Lys₆-Leu-enkephalin-Lys₆-OH);

Glu-His-Trp-Ser-Tyr-Gly-Leu-Arg-Pro-Gly-(Lys)₆-OH (GnRH-Lys₆-OH);

30

Glu-His-Trp-Ser-Tyr-Gly-Leu-Arg-Pro-Gly-(Lys-Glu)₃-OH (GnRH-(Lys-Glu)₃-OH); and

066720 0657450

H-Ser-Val-Ser-Glu-Ile-Gln-Leu-Met-His-Asn-Leu-Gly-Lys-His-Leu-Asn-Ser-Met-Glu-Arg-Val-Glu-Trp-Leu-Arg-Lys-Lys-Leu-Gln-Asp-Val-His-Asn-Phe-(Lys-Glu)₃-OH (PTH 1-34 human-(Lys-Glu)₃-OH).

5

26. A method for the preparation of a pharmacologically active peptide conjugate (X-Z) as defined in claim 2, comprising the steps of:

a) coupling an N- α -protected amino acid or N- α -protected dipeptide in the carboxyl
10 activated form, in the C-terminal activated form to an immobilised peptide sequence H-Z-SSM, thereby forming an immobilised N- α -protected peptide fragment,

b) removing the N- α -protecting group, thereby forming an immobilised peptide fragment having an unprotected N-terminal end,

15

c) coupling an additional N- α -protected amino acid in the carboxyl activated form, or an additional N- α -protected dipeptide in the C-terminal activated form to the N-terminal end of the immobilised peptide fragment, and repeating the removal/coupling step procedure in step b) and c) until the desired peptide sequence X is obtained, and then

20

d) cleaving off the peptide conjugate from the solid support material.

27. A method for the preparation of a pharmacologically active peptide conjugate (Z-X) as defined in claim 3, comprising the steps of:

25

a) coupling an N- α -protected amino acid, or an N- α -protected dipeptide to a solid support material (SSM), thereby forming an immobilised N- α -protected amino acid,

b) removing the N- α -protecting group, thereby forming an immobilised amino acid or
30 peptide fragment having an unprotected N-terminal end,

09341590.071399
062720 05574660

9/73

c) coupling an additional N- α -protected amino acid in the carboxyl activated form, or an additional N- α -protected dipeptide in the C-terminal activated form to the N-terminal end of the immobilised amino acid or peptide fragment, and repeating the removal/coupling step procedure in step b) and c) until the desired peptide sequence X is obtained,

5

d) coupling an additional N- α -protected amino acid in the carboxyl activated form, or an additional N- α -protected dipeptide in the C-terminal activated form to the N-terminal end of the immobilised peptide fragment, and repeating the removal/coupling step procedure in step b) and d) until the desired peptide sequence Z is obtained, and then

10

e) cleaving off the peptide conjugate from the solid support material.

28. A method for the preparation of a pharmacologically active peptide conjugate (Z-X-Z) as defined in claim 4, comprising the steps of:

15

a) coupling an N- α -protected amino acid in the carboxyl activated form, or an N- α -protected dipeptide in the C-terminal activated form to an immobilised peptide sequence H-Z-SSM, thereby forming an immobilised N- α -protected peptide fragment,

20 b) removing the N- α -protecting group, thereby forming an immobilised peptide fragment having an unprotected N-terminal end,

c) coupling an additional N- α -protected amino acid in the carboxyl activated form, or an additional N- α -protected dipeptide in the C-terminal activated form to the N-terminal end of
25 the immobilised peptide fragment, and repeating the removal/coupling step procedure in step b) and c) until the desired peptide sequence X is obtained, and then

d) coupling an additional N- α -protected amino acid in the carboxyl activated form, or an additional N- α -protected dipeptide in the C-terminal activated form to the N-terminal end of
30 the immobilised peptide fragment, and repeating the removal/coupling step procedure in step b) and d) until the desired peptide sequence Z is obtained, and then

e) cleaving off the peptide conjugate from the solid support material.

- Sub D³
29. A method for producing the peptide conjugate of claim 1, comprising
- a) introducing a nucleic acid sequence encoding said conjugate into a host cell;
 - 5 b) culturing said host cell and
 - c) isolating said conjugate from the culture.

30. A method for producing the peptide conjugate of claim 1, comprising
- a) culturing a recombinant host cell comprising a nucleic acid sequence encoding said
 - 10 conjugate under conditions permitting the production of said conjugate; and
 - b) isolating said conjugate from the culture.

a

31. The method according to claim 29 ~~or claim 30~~, wherein the nucleic acid sequence encoding said conjugate is contained within a nucleic acid construct or a vector.

15

32. A composition comprising a pharmacologically active peptide conjugate ~~as defined in any of the claims 1-25~~, and a pharmaceutical acceptable carrier.

according to claim 1

~~33. A composition comprising~~

- 20 H-Trp-Ala-Gly-Gly-Asp-Ala-Ser-Gly-Glu-(Lys-Glu)₃-OH,
 H-Trp-Ala-Gly-Gly-Asp-Ala-Ser-Gly-Glu-(Glu)₆-OH,
 H-Tyr-Gly-Gly-Phe-Leu-(Glu)₆-OH or
 H-Tyr-Gly-Gly-Phe-Leu-(Lys)₆-OH, and a pharmaceutical acceptable carrier.

25 34. A pharmacologically active peptide conjugate as defined in any of claims 1-26 for use in therapy.

35. A pharmacologically active peptide conjugate selected from the group consisting of
- H-Trp-Ala-Gly-Gly-Asp-Ala-Ser-Gly-Glu-(Lys-Glu)₃-OH,
 - 30 H-Trp-Ala-Gly-Gly-Asp-Ala-Ser-Gly-Glu-(Glu)₆-OH,
 - H-Tyr-Gly-Gly-Phe-Leu-(Glu)₆-OH and
 - H-Tyr-Gly-Gly-Phe-Leu-(Lys)₆-OH,

09341590-071399

for use in therapy.

36. Use of a pharmacologically active peptide conjugate as defined in any of claims 1-25 for the manufacture of a pharmaceutical composition for use in treatment of pain, HIV, 5 cancer, diabetes, incontinence, hypertension, amnesia, Alzheimer's disease, fever, depression, sex-hormone regulation, eating, schizophrenia, osteoporosis and insomnia.

Sub C6
37. A method for inhibiting neurons from transmitting pain impulses to the spinal cord, comprising administering to a subject in need thereof a conjugate comprising enkephalin 10 and a stabilising sequence, Z of 4-20 amino acids covalently attached to X, wherein Z is selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I



15

wherein R¹ and R² are selected from the group consisting of hydrogen, C₁₋₆-alkyl, phenyl, and phenyl-methyl, wherein C₁₋₆-alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents 20 selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R¹ and R² together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring, e.g. 2,4-diaminobutanoic acid and 2,3-diaminopropanoic acid;

25 wherein the ratio between the half-life of said peptide conjugate and the half-life of enkephalin when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least about 2, preferably at least about 3, such as at least about 5, more preferably at least about 7, such as at least about 9, e.g. at least about 10 or when X is inot orally absorbed said 30 peptide conjugate is orally absorbed in an amount effective to inhibit neurons from transmitting pain impulses.

094150.07139

38. Use of a conjugate comprising enkephalin and a stabilising sequence, Z of 4-20 amino acids covalently attached to X, wherein Z is selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I

5



wherein R¹ and R² are selected from the group consisting of hydrogen, C₁₋₆-alkyl, phenyl, and phenyl-methyl, wherein C₁₋₆-alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R¹ and R² together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring, e.g. 2,4-diaminobutanoic acid and 2,3-diaminopropanoic acid; and

Wherein the ratio between the half-life of said peptide conjugate and the half-life of enkephalin when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least about 2, preferably at least about 3, such as at least about 5, more preferably at least about 7, such as at least about 9, e.g. at least about 10 or when X is inot orally absorbed said peptide conjugate is orally absorbed for the manufacture of a pharmaceutical composition for use in treatment of pain.

39. A method for stimulating the release of growth hormone from the pituitary comprising administering to a subject in need thereof a conjugate comprising growth hormone releasing hormone or growth hormone releasing peptide and a stabilising sequence, Z of 4-20 amino acids covalently attached to X, wherein Z is selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I



0551420 20479PC1

wherein R^1 and R^2 are selected from the group consisting of hydrogen, C_{1-6} -alkyl, phenyl, and phenyl-methyl, wherein C_{1-6} -alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and
 5 phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C_{1-6} -alkyl, C_{2-6} -alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R^1 and R^2 together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring, e.g. 2,4-diaminobutanoic acid and 2,3-diaminopropanoic acid; and

10

Wherein the ratio between the half-life of said peptide conjugate and the half-life of growth hormone releasing hormone or growth hormone releasing peptide when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least about 2, preferably at least
 15 about 3, such as at least about 5, more preferably at least about 7, such as at least about 9, e.g. at least about 10 or when X is not orally absorbed said peptide conjugate is orally absorbed in an amount effective to stimulate the release of growth hormone.

~~40. Use of a conjugate comprising growth hormone releasing hormone or growth hormone releasing peptide and a stabilising sequence, Z of 4-20 amino acids covalently attached to X, wherein Z is selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I~~



25

wherein R^1 and R^2 are selected from the group consisting of hydrogen, C_{1-6} -alkyl, phenyl, and phenyl-methyl, wherein C_{1-6} -alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents
 30 selected from C_{1-6} -alkyl, C_{2-6} -alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R^1 and R^2 together with the carbon atom to which they are bound form a

054450-010574E50

cyclopentyl, cyclohexyl, or cycloheptyl ring, e.g. 2,4-diaminobutanoic acid and 2,3-diaminopropanoic acid; and

wherein the ratio between the half-life of said peptide conjugate and the half-life of growth hormone releasing hormone or growth hormone releasing peptide when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least about 2, preferably at least about 3, such as at least about 5, more preferably at least about 7, such as at least about 9, e.g. at least about 10 or when X is not orally absorbed, said peptide conjugate is orally absorbed, for the manufacture of a pharmaceutical composition for use in stimulating the release of growth hormone

41. A method for increasing hemoglobin levels comprising administering to a subject in need thereof a conjugate comprising EMP-1 and a stabilising sequence, Z of 4-20 amino acids covalently attached to X, wherein Z is selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I




wherein R¹ and R² are selected from the group consisting of hydrogen, C₁₋₆-alkyl, phenyl, and phenyl-methyl, wherein C₁₋₆-alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R¹ and R² together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring, e.g. 2,4-diaminobutanoic acid and 2,3-diaminopropanoic acid; and

- Wherein the ratio between the half-life of said peptide conjugate and the half-life of EMP-1 when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least

065720-0657460

about 2, preferably at least about 3, such as at least about 5, more preferably at least about 7, such as at least about 9, e.g. at least about 10 or when X is not orally absorbed said peptide conjugate is orally absorbed in an amount effective to increase hemoglobin levels.

5 42. Use of a conjugate comprising EMP-1 and a stabilising sequence, Z of 4-20 amino acids covalently attached to X, wherein Z is selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I



wherein R¹ and R² are selected from the group consisting of hydrogen, C₁₋₆-alkyl, phenyl, and phenyl-methyl, wherein C₁₋₆-alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and
15 phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R¹ and R² together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring, e.g. 2,4-diaminobutanoic acid and 2,3-diaminopropanoic acid;

20 wherein the ratio between the half-life of said peptide conjugate and the half-life of EMP-1 when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least about 2, preferably at least about 3, such as at least about 5, more preferably at least about 7,
25 such as at least about 9, e.g. at least about 10 or when X is not orally absorbed said peptide conjugate is orally absorbed for the manufacture of a pharmaceutical composition for use in treating anemia-by-increasing-hemoglobin-levels.

43. A method for preventing or treating bone loss by altering the balance between
30 osteoclastic and osteoblast activity comprising administering to a patient in need thereof a conjugate comprising parathyroid hormone and a stabilising sequence, Z of 4-20 amino acids covalently attached to X, wherein Z is selected from the group consisting of Ala, Leu,

662720" 06544660

Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I



5

wherein R¹ and R² are selected from the group consisting of hydrogen, C₁₋₆-alkyl, phenyl, and phenyl-methyl, wherein C₁₋₆-alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents
10 selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R¹ and R² together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring, e.g. 2,4-diaminobutanoic acid and 2,3-diaminopropanoic acid; and

15 Wherein the ratio between the half-life of said peptide conjugate and the half-life of parathyroid hormone when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least about 2, preferably at least about 3, such as at least about 5, more preferably at least about 7, such as at least about 9, e.g. at least about 10 or when X is not orally absorbed
20 said peptide conjugate is orally absorbed in an amount effective to treat or prevent bone loss.

~~44. Use of a conjugate comprising parathyroid hormone and a stabilising sequence, Z of 4-20 amino acids covalently attached to X, wherein Z is selected from the group consisting of
25 Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I~~



30 wherein R¹ and R² are selected from the group consisting of hydrogen, C₁₋₆-alkyl, phenyl, and phenyl-methyl, wherein C₁₋₆-alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and

09341590-07369

phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R¹ and R² together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring, e.g. 2,4-diaminobutanoic acid and 2,3-diaminopropanoic acid; and

Wherein the ratio between the half-life of said peptide conjugate and the half-life of parathyroid hormone when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least about 2, preferably at least about 3, such as at least about 5, more preferably at least about 7, such as at least about 9, e.g. at least about 10, or when X is not orally absorbed said peptide conjugate is orally absorbed, for the manufacture of a pharmaceutical composition for use in preventing or treating osteoporosis.

45. A method for reducing blood glucose levels comprising administering to a subject in need thereof a conjugate comprising glucagon-like peptide-1 and a stabilising sequence, Z of 4-20 amino acids covalently attached to X, wherein Z is selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I



wherein R¹ and R² are selected from the group consisting of hydrogen, C₁₋₆-alkyl, phenyl, and phenyl-methyl, wherein C₁₋₆-alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R¹ and R² together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring, e.g. 2,4-diaminobutanoic acid and 2,3-diaminopropanoic acid; and

0941590-0729

wherein the ratio between the half-life of said peptide conjugate and the half-life of glucagon-like peptide-1 when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least about 2, preferably at least about 3, such as at least about 5, more preferably at least about 7, such as at least about 9, e.g. at least about 10 or when X is not orally absorbed said peptide conjugate is orally absorbed in an amount effective to reduce blood glucose levels.

46. Use of a conjugate comprising glucagon-like peptide-1 and a stabilising sequence, Z of 4-20 amino acids covalently attached to X, wherein Z is selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I




wherein R¹ and R² are selected from the group consisting of hydrogen, C₁₋₆-alkyl, phenyl, and phenyl-methyl, wherein C₁₋₆-alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R¹ and R² together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring, e.g. 2,4-diaminobutanoic acid and 2,3-diaminopropanoic acid; and

wherein the ratio between the half-life of said peptide conjugate and the half-life of glucagon-like peptide-1 when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least about 2, preferably at least about 3, such as at least about 5, more preferably at least about 7, such as at least about 9, e.g. at least about 10 or when X is not orally absorbed said peptide conjugate is orally absorbed, for the manufacture of a pharmaceutical composition for use in treatment of diabetes.

09341590.071399
666720"0674E60


47. A method for regulating production of sex hormones comprising administering to a subject in need thereof a conjugate comprising gonadotropin releasing hormone and a stabilising sequence, Z of 4-20 amino acids covalently attached to X, wherein Z is selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I



wherein R¹ and R² are selected from the group consisting of hydrogen, C₁₋₆-alkyl, phenyl, and phenyl-methyl, wherein C₁₋₆-alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R¹ and R² together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring, e.g. 2,4-diaminobutanoic acid and 2,3-diaminopropanoic acid; and

wherein the ratio between the half-life of said peptide conjugate and the half-life of gonadotropin when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least about 2, preferably at least about 3, such as at least about 5, more preferably at least about 7, such as at least about 9, e.g. at least about 10 or when X is not orally absorbed said peptide conjugate is orally absorbed in an amount effective to regulate production of sex hormones.


48. Use of a conjugate comprising gonadotropin-releasing hormone and a stabilising sequence, Z of 4-20 amino acids covalently attached to X, wherein Z is selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I



wherein R^1 and R^2 are selected from the group consisting of hydrogen, C_{1-6} -alkyl, phenyl, and phenyl-methyl, wherein C_{1-6} -alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C_{1-6} -alkyl, C_{2-6} -alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R^1 and R^2 together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring, e.g. 2,4-diaminobutanoic acid and 2,3-diaminopropanoic acid; and

wherein the ratio between the half-life of said peptide conjugate and the half-life of gonadotropin releasing hormone when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least about 2, preferably at least about 3, such as at least about 5, more preferably at least about 7, such as at least about 9, e.g. at least about 10 or when X is not orally absorbed said peptide conjugate is orally absorbed and the ratio between the half-life of said peptide conjugate for the manufacture of a pharmaceutical composition for use in regulating the level of sex hormones.

49. A method for treating sleep disorders comprising administering to a subject in need thereof a conjugate comprising delta sleep inducing peptide and a stabilising probe, Z of 4-20 amino acids covalently attached to X, wherein Z is selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I



wherein R^1 and R^2 are selected from the group consisting of hydrogen, C_{1-6} -alkyl, phenyl, and phenyl-methyl, wherein C_{1-6} -alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C_{1-6} -alkyl, C_{2-6} -alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R^1 and R^2 together with the carbon atom to which they are bound form a

0341590-0239

cyclopentyl, cyclohexyl, or cycloheptyl ring, e.g. 2,4-diaminobutanoic acid and 2,3-diaminopropanoic acid; and

wherein the ratio between the half-life of said peptide conjugate and the half-life of delta-sleep inducing peptide when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least about 2, preferably at least about 3, such as at least about 5, more preferably at least about 7, such as at least about 9, e.g. at least about 10, or when the pharmacologically active peptide X is not orally absorbed, the conjugate will be orally absorbed, in an amount effective to treat said sleep disorder.

50. Use of a conjugate comprising delta-sleep inducing peptide and a stabilising probe, Z of 4-20 amino acids covalently attached to X, wherein Z is selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I



wherein R¹ and R² are selected from the group consisting of hydrogen, C₁₋₆-alkyl, phenyl, and phenyl-methyl, wherein C₁₋₆-alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R¹ and R² together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring, e.g. 2,4-diaminobutanoic acid and 2,3-diaminopropanoic acid; and

wherein the ratio between the half-life of said peptide conjugate and the half-life of Substance P when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C and in serum or plasma is at least about 2, preferably at least about 3, such as at least about 5, more preferably at least about 7, such as at least about 9, e.g. at least about 10, or when the pharmacologically active

0341590-0139
652729-0651460

~~peptide X is not orally absorbed, the conjugate will be orally absorbed, for the manufacture of a pharmaceutical composition for use in treating a sleep disorder~~

5 51. Use of a stabilising peptide sequence (Z) of 4-20 amino acid units for the preparation of a pharmacologically active peptide conjugate as defined in any of claims 1-26,

each amino acid unit in said stabilising peptide sequence Z being independently selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His,
10 Met, Orn, or amino acid units of the general formula I



wherein R¹ and R² independently are selected from hydrogen, C₁₋₆-alkyl, phenyl, and
15 phenyl-methyl, wherein C₁₋₆-alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R¹ and R² together with the carbon atom to which they are bound form a
20 cyclopentyl, cyclohexyl, or cycloheptyl ring;
or a salt thereof.

Add C7
Add D^{no}
Add E19
Add F13

09341590.071399
662720"0514E60